

10/23/00
JC944 U.S. PTO

10-24-00

A

Docket Number		4-100-8314C/C1C1
FILING BY "EXPRESS MAIL" UNDER 37 CFR 1.10		
EL 402023243US		OCT 23 2000
Express Mail Label Number		Date of Deposit

JC841 U.S. PTO
09/694209
10/23/00

Address to: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

UTILITY PATENT APPLICATION TRANSMITTAL AND FEE SHEET

Transmitted herewith for filing under 37 CFR §1.53(b)(1) is a **continuation** of prior Application No. 09/469,536, filed December 22, 1999.

Applicant (or identifier): HAEBERLIN ET AL.

Title: ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS

Enclosed are:

1. ☒ Specification (Including Claims and Abstract) - 14 pages
 2. ☐ Drawings - sheets
 3. ☐ Declaration and Power of Attorney
 - a. ☐ Newly executed (original or copy)
 - b. ☒ Copy from a prior application (signed or with indication that original was signed)
 - i. ☐ Deletion of Inventors
Signed statement attached deleting inventor(s) named in the prior application
 4. ☒ Incorporation By Reference
The entire disclosure of the prior application, from which a copy of the Declaration and Power of Attorney is supplied under Box 3b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
 5. ☐ Microfiche Computer Program (appendix)
 6. ☐ Nucleotide and/or Amino Acid Sequence Submission
 - ☐ Computer Readable Copy
 - ☐ Paper Copy
 - ☐ Statement Verifying Identity of Above Copies
 7. ☒ Preliminary Amendment
 8. ☐ Assignment Papers (Cover Sheet & Document(s))
 9. ☐ English Translation of
 10. ☐ Information Disclosure Statement
 11. ☐ Certified Copy of Priority Document(s)
 12. ☒ Return Receipt Postcard
 13. ☐ Other:
- ☒ The right to elect an invention or species that is different from that elected in parent Application No. 09/469,536 in the event of a restriction or election of species requirement that is identical or substantially similar to that made in said parent application is hereby reserved.

Filing fee calculation:

- ☒ Before calculating the filing fee, please enter the enclosed Preliminary Amendment.
☐ Before calculating the filing fee, please cancel claims

Basic Filing Fee							\$	710
Multiple Dependent Claim Fee (\$ 270)							\$	
Foreign Language Surcharge (\$ 130)							\$	
	For	Number Filed		Number Extra		Rate		
Extra Claims	Total Claims	14	-20	0	x	\$ 18	=	\$
	Independent Claims	2	-3	0	x	\$ 80	=	\$
TOTAL FILING FEE							\$	710

- ☒ Please charge Deposit Account No. 19-0134 in the name of Novartis Corporation in the amount of \$710. An additional copy of this paper is enclosed. The Commissioner is hereby authorized to charge any additional fees under 37 CFR §1.16 and §1.17 which may be required in connection with this application, or credit any overpayment, to Deposit Account No. 19-0134 in the name of Novartis Corporation.


Please address all correspondence to the address associated with Customer No. 001095, which is currently:

Thomas Hoxie
 Novartis Corporation
 Patent and Trademark Dept.
 564 Morris Avenue
 Summit, NJ 07901-1027

Please direct all telephone calls to the undersigned at the number given below and all telefaxes to (908) 522-6955.

Respectfully submitted,

Date: 10/23/00


 Michael U. Lee
 Attorney for Applicants
 Reg. No. 35,240
 Tel. No. (908) 522-6794

36841 U.S. PRO
 09/694209
 10/23/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

IN RE APPLICATION OF

HAEBERLIN ET AL.

APPLICATION NO: UNKNOWN

FILED: HEREWITH

FOR: ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS

Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT

Sir:

Prior to examination of the above-referenced case, please amend the application as follows:

IN THE SPECIFICATION

Please insert after the title, -- This application is a continuation of U.S. application Ser. No. 09/469,536, filed December 22, 1999, which is a continuation of U.S. application Ser. No. 09/077,398, filed May 28, 1998, which is a 371 of PCT/EP97/01800, filed April, 10, 1997. --.

IN THE CLAIMS

Please cancel claims 2-7.

Please amend claim 1 as follows:

1. (Amended) A pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to prevent release mycophenolate in the stomach [upper part of the intestinal tract].

Please add claims 8- 20 as follows:

8. The composition of claim 1 wherein the composition has an enteric coating and said enteric coating comprises cellulose acetate phthalate and trimellitate, or methacrylic acid copolymers containing at least 40 % methacrylic acid, or hydroxypropyl methylcellulose phthalate.

9. The composition of claim 8 wherein said coating comprises methacrylic acid copolymers containing at least 40 % methacrylic acid.

10. The composition of claim 8 wherein said composition is produced in a tablet form.

11. The composition of claim 10, wherein said tablet has a hardness between 10 and 70 N.

12. The composition of claim 8 wherein said composition is produced in a granule or pellet form.

13. The composition of claim 12 wherein said granule or pellet is contained in a capsule.

14. The composition of claim 1 wherein said salt is a mono-sodium salt.

15. The composition of claim 14 wherein said salt is in crystalline form.

16. The composition of claim 1 wherein said composition comprises from about 50 mg to 1.5 g of a pharmaceutically acceptable mycophenolate salt.

17. A pharmaceutical composition comprising a mycophenolate mono-sodium salt, the composition being adapted to prevent release mycophenolate in the stomach, wherein said composition has an enteric coating and said enteric coating comprises cellulose acetate phthalate and trimellitate, or methacrylic acid copolymers containing at least 40 % methacrylic acid, or hydroxypropyl methylcellulose phthalate.

18. The composition of claim 17 wherein said mono-sodium salt is in crystalline form.

19. The composition of claim 17 wherein said composition is in a tablet form and said tablet form has a hardness between 10 and 70 N.

20. The composition of claim 17 wherein said composition comprises from about 50 mg to 1.5 g of a pharmaceutically acceptable mycophenolate salt.

REMARKS

Claim 1 has been amended, and claims 8-20 have been added. Claims 2-7 have been cancelled. Please enter the above amendments before calculating fees for and examining the present application.

Respectfully submitted,

Novartis Corporation
Patent and Trademark Dept.
564 Morris Avenue
Summit, NJ 07901-1027
(908) 522-6794



Michael U. Lee
Attorney for Applicants
Reg. No. 35,240

Date:

10/23/00

ENTERIC COATED PHARMACEUTICAL COMPOSITIONS

This invention relates to mycophenolic acid.

Mycophenolic acid, also referred to herein as MPA, was first isolated in 1896, and has been extensively investigated as a pharmaceutical of potential commercial interest. It is known to have anti-tumor, anti-viral, immunosuppressive, anti-psoriatic, and anti-inflammatory activity [see e.g. W.A. Lee et al, *Pharmaceutical Research* (1990), 7, p. 161 - 166 and references cited therein]. Publications have appeared on MPA as an anti-cancer agent by Lilly scientists, see e.g. M.J. Sweeney et al., *Cancer Research* (1972), 32, 1795-1802, and by ICI scientists, see e.g. GB 1,157,099 and 1,203,328 and as an immunosuppressant agent see e.g. A. Mitsui et al. *J. Antibiotics* (1969) 22, p. 358-363. In the above-mentioned article by W.A. Lee et al it is stated that attempts have been made to increase the bio-availability or specificity of MPA by making derivatives. The poor bioavailability of the acid was thought to be caused by undetermined factors such as drug complexation in the gastrointestinal lumen, a narrow absorption window, metabolism before absorption etc.. The preparation of the morpholinoethyl ester, also known as mycophenolate mofetil (sometimes referred to herein as MMF), was described which had considerably higher bioavailability than MPA (100% for MMF and 43% for MPA). This derivative has been recently introduced commercially as an immunosuppressant for the treatment or prevention of organ or tissue transplant rejection, at daily dosages of from about 200 mg to about 3 grams p.o., e.g. about 2 g p.o. Patient compliance with MMF is not ideal, inter alia, because of side-effects e.g. gastro-intestinal side effects, the origin of which is not known.

We have now found, after exhaustive testing, that mycophenolate salts when enteric coated or adapted to be released in the upper part of the intestines, e.g. in the duodenum, jejunum and/or ileum, are effective, well-tolerated, pharmaceuticals particularly for immunosuppressive indications especially for the treatment or prevention of organ, tissue

or cellular allograft or xenograft rejection, e.g. after transplant, or the treatment or prevention of immune-mediated diseases (autoimmune diseases) and have interesting bioavailability and stability characteristics. Moreover fewer unit dosage forms are required to be administered than for MMF, leading to easier administration.

5 The present invention provides in one aspect a pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract (hereinafter referred to as a composition of the invention). The composition may be adapted in any conventional manner, preferably with means adapted to prevent release of the mycophenolate in the stomach and to ensure release in the upper
10 part of the intestinal tract. In a further aspect the invention provides a pharmaceutical composition comprising a coated pharmaceutically acceptable mycophenolate salt.

Such salts are cationic salts, e.g. of alkali metals, especially the sodium salts. Sodium mycophenolate salts are known, e.g. in South African Patent 68/4959. We prefer to use the mono-sodium salt. This may be obtained in crystalline form by recrystallization from
15 acetone/ethanol if necessary with water; Mpt. 189 - 191°C.

The invention provides, more specifically, a solid enteric-coated composition in unit dose form for oral application, the core of the composition containing sodium mycophenolate in solid or liquid form.

The term "core" comprises sodium mycophenolate (or other cationic salt) if desired in
20 admixture with further physiologically acceptable material, that can be surrounded by an enteric-coating. The term "core" comprises, in a wide sense, not only tablets, pellets or granules but also capsules, e.g. soft or hard capsules of gelatine or starch. Such cores may be produced in conventional manner. We have found that the mycophenolate salts, particularly the sodium salt, are particularly interesting for the production of tablets. When
25 tablet cores are used they have preferably a hardness of from ca. 10 to 70 N.

The pellets or granules may, after application of the enteric-coating as described hereinafter may be used as such or to fill capsules, e.g. hard gelatine capsules. If desired the capsules may be alternatively enteric-coated, e.g. in conventional manner.

5 Other pharmaceutically acceptable ingredients may be present in the cores, e.g. those conventionally used in the preparation of pharmaceutically compositions, e.g. fillers, e.g. lactose, glidants, e.g. silica, and lubricants, e.g. magnesium stearate.

10 The term "enteric coating" comprises any pharmaceutically acceptable coating preventing the release of the active agent in the stomach and sufficiently disintegrating in the intestine tract (by contact with approximately neutral or alkaline intestine juices) to allow the resorption of the active agent through the walls of the intestinal tract. Various in vitro tests for determining whether or not a coating is classified as an enteric coating have been published in the pharmacopoeia of various countries.

15 More specifically, the term "enteric coating" as used herein refers to a coating which remains intact for at least 2 hours, in contact with artificial gastric juices such as HCl of pH 1 at 36 to 38°C and preferably thereafter disintegrates within 30 minutes in artificial intestinal juices such as a KH_2PO_4 buffered solution of pH 6.8.

The thickness of the coating may vary and depends inter alia on its permeability in water and acids. A typical coating may be about 16-30, e.g. 16-20 or to 25, mg on a size 1 gelatine capsule. Similar thicknesses may be applied in other formulations.

20 In general satisfactory results are obtained with a coating of 5 - 100 μm , preferably 20 - 80 μm thickness. The coating is suitably selected from macromolecular polymers. Suitable polymers are listed in e.g. L. Lachman et al. The Theory and Practice of Industrial Pharmacy, 3rd Ed, 1986, p. 365 - 373, H. Sucker et al, Pharmazeutische Technologie, Thieme, 1991, p. 355 - 359, Hagers Handbuch der pharmazeutischen Praxis, 4th Ed. Vol.
25 7, pages 739 to 742 and 766 to 778, (Springer Verlag, 1971) and Remington's Pharmaceutical Sciences, 13th Ed., pages 1689 to 1691 (Mack Publ., Co., 1970) and

comprise e.g. cellulose ester derivatives, cellulose ethers, acrylic resins, such as methylacrylate copolymers and copolymers of maleic acid and phthalic acid derivatives.

5 The preferred films are made from cellulose acetate phthalate and trimellitate; methacrylic acid copolymers, e.g. copolymers derived from methylacrylic acid and esters thereof, containing at least 40% methylacrylic acid; and especially hydroxypropyl methylcellulose phthalate.

Methylacrylates include those of molecular weight above 100,000 daltons based on, e.g. methylacrylate and methyl or ethyl methylacrylate in a ratio of about 1:1. Typical products include Endragit L, e.g. L 100-55, marketed by Rohm GmbH, Darmstadt, Germany.

10 Typical cellulose acetate phthalates have an acetyl content of 17-26% and a phthalate content of from 30-40% with a viscosity of ca. 45-90cP.

Typical cellulose acetate trimellitites have an acetyl content of 17-26%, a trimellityl content from 25 - 35 % with a viscosity of ca. 15-20 cS. An example of an appropriate cellulose acetate trimellitate is the marketed product CAT (Eastman Kodak Company, USA).

15 Hydroxypropyl methylcellulose phthalates, typically have a molecular weight of from 20,000 to 100,000 daltons e.g. 80,000 to 130,000 daltons, e.g. a hydroxypropyl content of from 5 to 10%, a methoxy content of from 18 to 24% and a phthalyl content from 21 to 35%.

20 An example of an appropriate cellulose acetate phthalate is the marketed product CAP (Eastman Kodak, Rochester N.Y., USA).

Examples of suitable hydroxypropyl methylcellulose phthalates are the marketed products having a hydroxypropyl content of from 6-10%, a methoxy content of from 20-24%, a phthalyl content of from 21-27%, a molecular weight of about 84,000 daltons known under the trade mark HP50 and available from Shin-Etsu Chemical Co. Ltd., Tokyo, Japan, and

having a hydroxypropyl content, a methoxyl content, and a phthalyl content of 5-9%, 18-22% and 27-35% respectively, and a molecular weight of 78,000 daltons, known under the trademark HP55 and available from the same supplier.

A preferred coating is HP 50.

- 5 The enteric coating may be carried out in conventional manner, e.g. so that the cores are sprayed with a solution of the enteric-coating.

Suitable solvents for the enteric-coating are for example organic solvents, e.g. an alcohol such as ethanol, a ketone such as acetone, halogenated hydrocarbons such as CH_2Cl_2 or mixtures of such solvents, e.g. ethanol /acetone, e.g. 1:1 to 10:1.

- 10 Conveniently a softener such as di-n-butylphthalate or triacetin is added to such a solution, e.g. in a ratio of coating material to softener of from 1: about 0.05 to about 0.3.

If desired for cellulose phthalates and other acidic coating materials an ammonium salt may be found and an aqueous solution may be used.

A fluidized bed coater may be used for coating.

- 15 Conveniently the cores are treated at room temperature or warmed up to 40°C e.g. by means of warm air of 40° up to 70°C , before spraying. To avoid a sticking of the cores the spray procedure is preferably interrupted at certain time intervals and the cores then warmed up again. It is, however, also possible to proceed without interruption of the spray procedure, e.g. by automatic regulation of the spray amount taking into account the temperature of
20 exhaust air and/or cores.

The spray pressure may vary within wide ranges, in general satisfactory results are obtained with a spray pressure of from about 1 to about 1.5 bar.

The compositions of the invention are useful as immunosuppressants as indicated by standard tests.

The activity and characteristics of the compositions of the invention may be indicated in standard

5 a) clinical trials, e.g. observing the first acute rejection episodes or treatment failure
six months after transplant of kidneys or maintaining a rejection - free state within
6 months after imitiation of treatment with the invention. The compositions of the
invention are administered at a dose in the range of 0.5 to 2.0 g/day e.g. about 1.5
g /day and decrease the acute rejection rates when administered during the period
10 around transplant surgery, and maintain a rejection-free state in patients who are 3
months or more after transplantation. Thus the compositions of the invention may
be administered during the initial 72 hours after transplantation at dose of about 0.5
g administered twice a day in combination with a conventional steroid and
cyclosporin, e.g. as NEORAL for which the cyclosporin dose is the conventional
15 dose e.g. ca. 8 ± 3 mg/kg for renal transplants. The steroid dose is to be
administered at about 2.5 mg /kg for 4 days after transplant, 1 mg/kg thereafter for
1 week, 0.6 mg/kg thereafter for 2 weeks thereafter 0.3 mg/kg for $\frac{1}{2}$ month for
prednisone.

and in

20 b) animal trials e.g. observing the kidney allograft reaction in rat. In this test one
kidney from a female fisher 344 rat is transplanted onto the renal vessel of a
unilaterally (left side) nephrectomized WF recipient rat using an end-to-end
anastomosis. Ureteric anastomosis is also end-to-end. Treatment commences on the
day of transplantation and is continued for 14 days. A contralateral nephrectomy is
25 done seven days after transplantation, leaving the recipient relying on the
performance of the donor kidney. Survival of the graft recipient is taken as the
parameter for a functional graft. Typical doses of the compositions of the invention

are from about 1 to 30 mg/kg p.o.

The compositions of the invention are particularly useful for the following conditions:

- a) Treatment and prevention of native or transgenic organ, tissue or cellular allograft or xenograft transplant rejection, e.g. for the treatment of recipients of e.g. heart, lung, combined heart-lung, liver, kidney, pancreatic, skin, pancreatic islet cell, neural cell or corneal transplant; including treatment and prevention of acute rejection; treatment and prevention of hyperacute rejection, e.g. as associated with xenograft rejection; and treatment and prevention of chronic rejection, e.g. as associated with graft-vessel disease. The compositions of the invention are also indicated for the treatment and prevention of graft-versus-host disease, such as following bone marrow transplantation.
- b) Treatment and prevention of autoimmune diseases, e.g. immune-mediated diseases and inflammatory conditions, in particular inflammatory conditions with an etiology including an immunological component such as arthritis (for example rheumatoid arthritis, arthritis chronica progrediente and arthritis deformans) and rheumatic diseases. Specific immune-mediated diseases for which the compositions of the invention may be employed include, autoimmune hematological disorders, including, but not limited to hemolytic anaemia, aplastic anaemia, pure red cell anaemia and idiopathic thrombocytopenia), systemic lupus erythematosus, polychondritis, sclerodoma, Wegener granulosis, dermatomyositis, polymyositis, chronic active hepatitis, primary biliary cirrhosis, myasthenia gravis, psoriasis, Steven-Johnson syndrome, pemphigus, idiopathic sprue, inflammatory bowel diseases (including e.g. ulcerative colitis and Crohn's disease), endocrine ophthalmopathy, Graves disease, sarcoidosis, multiple sclerosis, primary biliary cirrhosis, juvenile diabetes (diabetes mellitus type I), non-infectious uveitis (anterior and posterior), keratoconjunctivitis sicca and vernal keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis, vasculitides, glomerulonephritides (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy) and juvenile dermatomyositis.

Appropriate dosages of the compositions of the invention will of course vary, e.g. depending on the condition to be treated (for example the disease type or the nature of resistance), the MPA salt used, the effect desired and the mode of administration.

5 In general however satisfactory results are obtained on administration e.g. orally at dosages on the order of from about 1 to about 30 mg salt per kg animal body weight per day, administered once or in divided doses up to 4 times per day. Suitable daily dosages for patients are thus in the order of 200 mg to 3 g p.o. salt e.g. from about 50 to 100% that of mycophenolate mofetil. For the preferred mono sodium salt the dosage of the salt is about two thirds that of mycophenolate mofetil.

10 Representative unit dosage forms contain from about 50 mg, e.g. 100 mg, to about 1.5 g of the pharmaceutically acceptable mycophenolate salt.

The bioavailability characteristics of compositions of the invention may be determined in conventional manner, e.g. by oral administration to beagle dogs. Dosages are typically 50 mg salt animal e.g. ca 3- 5 mg salt /kg animal body weight. Dogs are adult (ca. 10 kg e.g. 15 6 - 14 kg) and fasted. Three hours after administration ca. 200 g food is administered. Blood samples are taken from the cephalic vein, before administration and 10, 30, and 45 minutes, 1, 1.5, 2, 3, 4, 6, 8, 12, and 24 hours, after administration. Plasma levels of free MPA are determined by HPLC analysis (with UV detection).

20 In a relative bioavailability trial as described above in male beagle dogs dosages of 3.8 mg salt/kg animal body weight p.o. were administered with the Example 1 composition as described hereinafter and with a MPA or MMF formulation corresponding to the Example 1 composition but containing an identical amount of MPA or commercially available MMF.

Results are as follows:-

	MPA (AUC Relative Bioavailability, Frel [ng.hr.ml ⁻¹]	Ex 1	MPA	MMF
5	Mean	4612 (218)	3579 (174)	2709 (100)
	Median	4204 (168)	2911 (182)	2513 (100)
	SD	939	1889	1363
	CV	20	53	50
10	Cmax [ng/ml] (Relative Cmax)			
	Mean	5391 (313)	3683 (227)	2052 (100)
	Median	5359 (367)	2719 (172)	1462 (100)
	SD	1847	2504	945
	CV (%)	34 (46)	68 (87)	46 (0)
15	The coefficients of variation (CV) of AUC (20%) and Cmax (34%) of the Example 1 composition are significantly less than those of the reference compositions, indicating less inter-subject and intra-subject variability with the Example 1 composition.			
	The area under the curve (AUC) and Cmax with the Example 1 composition are higher than those of the reference compositions.			
20	Naturally the advantageous bioavailability characteristics of the present compositions may be ascertained in standard clinical bioavailability trials. For example, doses from 200 mg to 1.5 g of the Example 1 composition and MPA, and MMF may be administered to 12 healthy volunteers in single doses in a cross-over trial. Increased AUC and C _{max} may be observed for the Example 1 composition.			

The compositions of the present invention are surprisingly tolerated better than MMF, inducing less gastro-intestinal side effects such as diarrhoea and burning. They show less long term side effects e.g. in the colon.

5 The compositions of the invention may be administered as the sole active ingredient or with another immunosuppressant e.g. together with simultaneous or separate administration of other immunosuppressants, for example, in immunosuppressive applications such as prevention and treatment of graft vs. host disease, transplant rejection, or immune-mediated disease, the compositions of the invention may be used in combination with cyclosporins or ascomycins, or their immunosuppressive analogs, e.g. cyclosporin A, FK- 506
10 (tacrolimus), etc., rapamycin; corticosteroids; cyclophosphamide; azathioprine; methotrexate; brequinar; leflunomide; mizoribine; deoxyspergualin; analogues thereof, and immunosuppressive monoclonal antibodies, e.g., monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, CTLA4, B7, CD45, or CS58 or their ligands; or other immunomodulatory compounds.

15 When the compositions of the invention are co-administered with such other immunosuppressants the dosages of the other immunosuppressants may be reduced e.g. to one-half to one-third their dosages when used alone.

Representative doses for ciclosporin to be used are e.g. 1 to 10, e.g. 1 to 2 mg/kg/day.

20 The present invention provides in another aspect the use, method and compositions as defined hereinafter in the claims.

Insofar as details of excipients are not described herein, these are known, or available e.g. in the Handbook of Pharmaceutical Excipients, Second Edition, edited by Ainley Wade and Paul J. Weller, American Pharmaceutical Association, Washington, USA and Pharmaceutical Press, London; and Lexikon der Hilfsstoffe für Pharmazie, Kosmetik and
25 angrenzende Gebiete edited by H.P. Fiedler, 4th Edition, Editio Cantor, Aulendorf and earlier editions.

Following is a description by way of example only of compositions of this invention:

EXAMPLE 1:

COMPOSITION

Capsule contents

5	MPA mono sodium salt	53.43 mg (= 50 mg MPA)
	Lactose	256.57 mg
	(1:1 mixture of 100/200 mesh)	
	Silica (Aerosil)	3.1 mg
	Magnesium stearate	1.55 mg
10		<hr/>
		314.65 mg

Capsule is size 1

Enteric coating (ca 17 mg)

15	Hydroxypropyl methyl cellulose	9 parts
	phthalate (HP50)	
	Triacetin	1 part

PROCEDURE

20 The capsule ingredients are mixed and filled into size 1 capsules. The capsules are coated in a fluidized bed coater with a solution of the enteric coating ingredients in ethanol (containing 10% acetone). The coating on each capsule is about 17 mg. The capsules meet the enteric coating test described herein and do not disintegrate within 2 hours in artificial gastric juices (pH 1, HCl). The compositions are stable, e.g for 2 years at room temperature,

If desired larger capsules containing 534.3 mg MFA mono sodium salt may be made in analogous manner, reducing the amount of lactose. These are well tolerated in clinical trials.

EXAMPLE 2:

5 Capsules of size 1 are made up as in Example 1. A solution for enteric coating was made up as follows:

Hydroxypropyl methyl cellulose phthalate (HP50)	270 g
Triacetin	30 g
Acetone	900 g
10 Ethanol	1800 g

600 g of this enteric coating solution was used for 1 kg of capsules (ca. 2400). The amount of coating applied to each capsule was about 25 mg giving a film thickness of 5-6 mg/cm².

CLAIMS:

1. A pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract.
2. A pharmaceutical composition comprising an enteric coated pharmaceutically acceptable mycophenolate salt.
3. Use of an enteric coated pharmaceutically acceptable mycophenolate salt or composition of claim 1 or 2 in the preparation of a medicament for immunosuppression, particularly for prevention or treatment of native or transgenic organ, tissue or cellular allograft or xenograft transplant rejection; for the treatment or prevention of immune-mediated and/or inflammatory diseases; optionally with the simultaneous or separate administration of another immunosuppressant.
4. A method of immunosuppressing a subject which comprises administering a therapeutically effective amount of enteric coated pharmaceutically acceptable mycophenolate or a composition of claim 1 or 2 to a subject in need of such immunosuppression, optionally with the simultaneous or separate administration of another immunosuppressant.
5. A composition containing an enteric coated pharmaceutically acceptable mycophenolate salt or a composition of claim 1 or 2 and another immunosuppressant for simultaneous, sequential or separate administration.
6. A composition, use or method according to any preceding claim wherein the sodium salt is the mono-sodium salt.
7. A composition, use or method according to any preceding claim wherein another immunosuppressant is present, e.g. ciclosporin.

ENTERIC COATED PHARMACEUTICAL COMPOSITIONS

Abstract of the Disclosure

5 This invention provides a pharmaceutical composition comprising a mycophenolate salt, the composition being adapted to release mycophenolate in the upper part of the intestinal tract.

6300/RR/ME

20-Feb-1997

m: spec \ 100 \ 8314.wpc

US Case 4 -100-8314/A

DECLARATION AND POWER OF ATTORNEY FOR UNITED STATES PATENT APPLICATION

☒ Original

☐ Supplemental

☐ Substitute

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, and

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if more than one name is listed below) of the subject matter which is claimed and for which a United States patent is sought on the invention entitled

ENTERIC-COATED PHARMACEUTICAL COMPOSITIONS

the specification of which:

☐ is attached hereto.

☐ was filed on _____ as Application No. _____
(day/month/year)

and, if this box (☐) contains an *

☐ was amended on _____
(day/month/year)

☒ was filed as Patent Cooperation Treaty international Application No.

PCT/EP 97/01800 on 10/04/97
(day/month/year)

and, if this box (☐) contains an *

☐ entered the national stage in the United States and was accorded Application No. _____

and, if this box (☐) contains an *

☐ was amended, subsequent to entry into the national stage, on _____
(day/month/year)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above and, if this application was filed as a Patent Cooperation Treaty international application, by any amendments made during the international stage (including any made under Patent Cooperation Treaty Rule 91, Article 19 and Article 34).

I acknowledge my duty to disclose all information which is known by me to be material to the patentability of this application as defined in 37 C.F.R. § 1.56.

I hereby claim the benefit under 35 U.S.C. §119(a)-(d) or §365(b) of any foreign application(s) for patent or inventor's certificate listed below and under 35 U.S.C. §365(a) of any Patent Cooperation Treaty international application(s) designating at least one country other than the United States listed below and have also listed below any foreign application(s) for patent or inventor's certificate and Patent Cooperation Treaty international application(s) designating at least one country other than the United States for the same subject matter and having a filing date before that of the application the priority of which is claimed for that subject matter:

COUNTRY/REGION (OR P.C.T.)	APPLICATION No.	FILING DATE (day/month/year)	PRIORITY CLAIMED	
GB	9607564.3	12/04/96	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
GB	9622028.0	24/10/96	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
			<input type="checkbox"/> Yes	<input type="checkbox"/> No
			<input type="checkbox"/> Yes	<input type="checkbox"/> No
			<input type="checkbox"/> Yes	<input type="checkbox"/> No

I hereby claim the benefit under 35 U.S.C. § 119 (e) of any United States provisional application(s) listed below:

APPLICATION NO.	FILING DATE (day/month/year)
-----------------	---------------------------------

I hereby claim the benefit under 35 U.S.C. §120 of any United States application(s) listed below and under 35 U.S.C. §365(c) of any Patent Cooperation Treaty international application(s) designating the United States listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in said prior application(s) in the manner required by the first paragraph of 35 U.S.C. §112, I acknowledge my duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. §1.56 which became available between the filing date(s) of the prior application(s) and the national or Patent Cooperation Treaty international filing date of this application:

United States Application No.	United States Filing Date (day/month/year)	Status (Pending, Abandoned or U.S. Patent No.)	International Application No. and Filing Date
----------------------------------	--	--	--

I hereby appoint the registered practitioners associated with Customer No. 001095, respectively and individually, as my attorneys and agents, with full power of substitution and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

If these brackets contain an X [X], I hereby authorize the registered practitioners associated with Customer No. 001095 and any others acting on my behalf to take any action relating to this application based on communications from the Patents and Trademarks Division of Novartis Services AG, Basle, Switzerland, or an affiliate thereof or a successor thereto, without direct communication from me.

Please address all communications to Michael W. Glynn, Novartis Corporation, Patent and Trademark Department, 564 Morris Avenue, Summit, NJ 07901-1027.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full name of sole
or first joint inventor

Barbara HAEBERLIN

Inventor's signature

Barbara Haebelin

Date

13-05-98
(day/month/year)

Residence

4125 Riehen, Switzerland

Citizenship

Swiss

Post Office Address

Haselrain 77
4125 Riehen
Switzerland

IMPORTANT: Before this declaration is signed, the patent application (the specification, the claims and this declaration) must be read and understood by each person signing it, and no changes may be made in the application after this declaration has been signed.

Full name of second
joint inventor, if any

Ching-Pong MAK

Inventor's signature

Ching Pong Mak

Date

May 18 98
(day/month/year)

Residence

4106 Therwil, Switzerland

Citizenship

British

Post Office Address

Ob dem Fichtenrain 20
4106 Therwil
Switzerland

Full name of third
joint inventor, if any

Armin MEINZER

Inventor's signature

Armin Meiner

Date

14-05-98
(day/month/year)

Residence

79426 Buggingen, Germany

Citizenship

German

Post Office Address

Fliederweg 2
79426 Buggingen
Germany

Full name of fourth
joint inventor, if any

Jacky VONDERSCHER

Inventor's signature

Jacky Vonderscher

Date

May 20, 98
(day/month/year)

Residence

68400 Riedisheim, France

Citizenship

French

Post Office Address

29, rue de Bruebach
68400 Riedisheim
France